

Available online at www.sciencedirect.com**ScienceDirect**journal homepage: www.elsevier.com/locate/ajps

Stability study of resveratrol-loaded emulsions using pectin as an emulsifier

Porntip Benjasirimongkol^{a,b,*}, Pornsak Sriamornsak^{a,b}

^a Department of Pharmaceutical Technology, Silpakorn University, Nakhon Pathom 73000, Thailand

^b Pharmaceutical Biopolymer Group (PBiG), Faculty of Pharmacy, Silpakorn University, Nakhon Pathom 73000, Thailand

ARTICLE INFO

Article history:

Available online 25 November 2015

Keywords:

Resveratrol

Pectin

Emulsion stability

Homogenization

Phenolic compound such as resveratrol has been suggested to prevent cancer and heart diseases because of its anti-oxidant and anti-inflammatory activities [1]. However, therapeutic application of resveratrol is quite limited due to its poor solubility, low bioavailability, photosensitive and auto-oxidation properties [2]. The aim of this study was to prepare resveratrol-loaded emulsions using pectin as an emulsifier. The emulsions should be stable for up to 7 days, before further manufacturing processes.

Resveratrol-loaded emulsions were prepared by simple homogenization using Imwitor® 742 as an oil phase (10% w/w). Resveratrol was dissolved in Imwitor® 742; solubility of resveratrol in Imwitor® 742 was 9.53 ± 0.52 mg/mL. The influence of type (i.e., low methoxy pectin (LMP) and high methoxy pectin (HMP)) and concentration (i.e., 2% and 3% w/w) of pectin on the droplet size, zeta potential, viscosity, stability (by visually observed percent creaming and automatic stability analysis) and morphology of emulsions were examined. The stability of emulsions after storage at ambient temperature (25 °C) for 7 days and under environmental stress condition (i.e.,

centrifugation test) was compared with the freshly prepared emulsions (day 0). The microscopic image of emulsions containing HMP (3% w/w) and resveratrol is shown in Fig. 1A. The emulsion droplet size was increased from 19.34 ± 2.6 μm (day 0) to 26.95 ± 1.76 μm (day 7). From visual observation, emulsions containing HMP (3% w/w) and resveratrol did not show phase separation (100% creaming) after centrifugation test and 7-day stability test. However, the results by automatic stability analyzer showed the difference between the emulsions after centrifugation test and 7-day stability test (Fig. 1B). It could be concluded that the emulsions containing HMP (3% w/w) and resveratrol provide the most stable emulsions compared to other formulations.

Acknowledgment

The authors wish to acknowledge Herbstreith & Fox KG (Germany) who kindly donate pectin samples.

* E-mail address: benjasirimongko_p@silpakorn.edu.

Peer review under responsibility of Shenyang Pharmaceutical University.

<http://dx.doi.org/10.1016/j.ajps.2015.11.037>

1818-0876/© 2016 Production and hosting by Elsevier B.V. on behalf of Shenyang Pharmaceutical University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1 – (A) Microscopic image and (B) backscattering patterns of emulsions containing HMP (3% w/w) and resveratrol.

REFERENCES

- [1] Pace-Asciak CR, Hahn S, Diamandis EP, et al. The red wine phenolics trans-resveratrol and quercetin block human platelet aggregation and eicosanoid synthesis: implications for protection against coronary heart disease. *Clin Chim Acta* 1995;235:207–219.
- [2] Amri A, Chaumeil JC, Sfar S, et al. Administration of resveratrol: what formulation solutions to bioavailability limitations? *J Control Release* 2012;158:182–193.